



Dear Parent/Guardian:

As your 11th or 12th grader prepares for the next phase of life, make sure he/she has all the vaccines that are recommended at this age. Many colleges, universities, and employers require vaccinations. That is why we are starting a program called “Vaccinate Before You Graduate (VBYG).”

To make things easier for your family, Howard County Public Schools is working with the Maryland Partnership for Prevention and Howard County Health Department to hold VBYG immunization clinics at your child’s school. This is the same partnership that provides flu vaccinations in the fall. These clinics for 11th and 12th graders will be held in April.

The immunizations will be given at **no cost to you. You will NOT be charged a deductible or copay. You do NOT have to be present** and you do not need insurance to be vaccinated.

Graduating Seniors Might Need...		
Vaccine Name	Why is it Needed?	When is it Needed?
Meningococcal Meningitis (MCV4) Booster	Meningitis is very contagious and can be deadly. Prevents four types of meningitis.	First dose at 11 or 12 years old. Second dose recommended five years after the first dose.
Meningococcal Meningitis Type B (MenB) Vaccine	It is responsible for most college outbreaks of meningitis.	First vaccination at about 16 years old. The second one is needed at least six months later.
Human Papilloma Virus (HPV) Vaccine	Prevents nine types of cancer. One of only two vaccines that can prevent cancer.	For males and females 9 to 26 years old. Depending on age, two or three vaccinations needed at least six months apart.
Tdap Vaccine	Prevents tetanus, diphtheria, and pertussis.	Now, if you didn’t get one after 11 years old.
When you fill out the consent form, initial beside all vaccines you would like your child to receive. We will check school records and the Maryland immunization registry, ImmuNet, see which ones are needed.		

If you want your child to be vaccinated, please:

1. Go to www.vaccineconsent.com and fill out a consent form online.

OR

2. Fill out every section of the Consent Form **on the back of this letter**. Write neatly in ink.
3. If you have insurance, use your insurance card to fill out your insurance information accurately.
4. Read the Vaccine Information Statement(s) at www.immunize.org/vis/ and see your doctor or school nurse with any questions.
5. **Return the consent form to your child’s school by March 22, 2019.**

We will send a record home with your child on the day he or she has been vaccinated. If your child does not get vaccinated on clinic day, we will send a note letting you know why.

We are pleased to provide this service to make sure your child enters the next phase of life protected from these serious diseases. Please call your child’s health care provider if you have any questions.

Howard County Consent for 2019 Recommended Vaccinations for 11th and 12th Graders

Please Print Clearly in Ink

Student's LAST NAME	Student's FIRST NAME	MI	Student's Date of Birth	Age	Sex	Grade
Parent/Guardian LAST NAME	FIRST NAME	MI	____/____/____		F M	
Address			Cell/Home Phone		Email Address	
City		ZIP Code	School Name			

HEALTH INSURANCE INFORMATION – PLEASE FILL OUT COMPLETELY AND ACCURATELY

Please copy this information from YOUR INSURANCE CARD. We will bill your insurance. **You will NOT be charged** a co-pay or a deductible.

Type of Insurance:		<input type="checkbox"/> Private Insurance or Medical Assistance	<input type="checkbox"/> My child does not have health insurance <small>(Your child will not be turned away because of no insurance)</small>																				
Insurance Company Name	Member ID Number (write in boxes below)																						
	<table border="1" style="width: 100%; height: 20px; border-collapse: collapse;"> <tr> <td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td> </tr> </table>																						
	Group Number (write in boxes below)																						
	<table border="1" style="width: 100%; height: 20px; border-collapse: collapse;"> <tr> <td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td><td style="width: 10%;"></td> </tr> </table>																						
FOR PRIVATE INSURANCE ONLY.																							
Policy Holder's/Insured Adult's Name	Relationship to Student	Insured Adult's Birthdate	Any Other # from Insurance Card																				

1. Do any of the following apply to your child? (If you answer YES to any question, your child might not be vaccinated.)

Yes	No		Yes	No	
<input type="checkbox"/>	<input type="checkbox"/>	Has had a serious reaction to a vaccine in the past?	<input type="checkbox"/>	<input type="checkbox"/>	Has had Guillain-Barre syndrome?
<input type="checkbox"/>	<input type="checkbox"/>	Has an allergy to a component in a vaccine?	<input type="checkbox"/>	<input type="checkbox"/>	Has a serious allergy to anything? List _____
<input type="checkbox"/>	<input type="checkbox"/>	Has brain or nervous system problems?	<input type="checkbox"/>	<input type="checkbox"/>	For females: Is pregnant?

PLEASE INITIAL BESIDE THE VACCINE(S) you would like your child to receive. We will check to be sure it is needed.

Parent's Initials	Vaccine Name	VIS Date
	Tdap	2/24/15
	Meningococcal ACWY (MCV4)	8/24/18
	Meningitis B (MenB)	8/9/2016
	Human Papilloma Virus (HPV)	12/2/16

**FOR OFFICE USE ONLY
PARENTS: DO NOT WRITE IN THIS SPACE**

Manufacturer	Site	Route	Lot Number
		IM	
		IM	
		IM	
		IM	

CONSENT FOR VACCINATION(S) - YOU MUST SIGN HERE FOR YOUR CHILD TO BE VACCINATED

By signing this form, I give permission for my child to be vaccinated with the vaccines listed above, my insurance company to be billed, and vaccine(s) entered into ImmuNet, Maryland's immunization registry. Further, I agree that the information above is correct, and:

- (1) I have read the current Vaccine Information Statement for each vaccine(s) or someone has read it to me;
- (2) I understand the risks and benefits of getting the vaccine(s) I have consented for my child to receive; and
- (3) Any questions I had about the vaccine(s) have been answered.

Signature of Parent/Legal Guardian _____ Date: ____/____/____

OFFICE USE ONLY – PARENTS: DO NOT WRITE IN THIS SPACE

Date VIS Given/Vaccine Administered	PRINT Name of Vaccine Administrator	Notes

Meningococcal ACWY Vaccine: What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Meningococcal disease is a serious illness caused by a type of bacteria called *Neisseria meningitidis*. It can lead to meningitis (infection of the lining of the brain and spinal cord) and infections of the blood. Meningococcal disease often occurs without warning—even among people who are otherwise healthy.

Meningococcal disease can spread from person to person through close contact (coughing or kissing) or lengthy contact, especially among people living in the same household.

There are at least 12 types of *N. meningitidis*, called “serogroups.” Serogroups A, B, C, W, and Y cause most meningococcal disease.

Anyone can get meningococcal disease but certain people are at increased risk, including:

- Infants younger than one year old
- Adolescents and young adults 16 through 23 years old
- People with certain medical conditions that affect the immune system
- Microbiologists who routinely work with isolates of *N. meningitidis*
- People at risk because of an outbreak in their community

Even when it is treated, meningococcal disease kills 10 to 15 infected people out of 100. And of those who survive, about 10 to 20 out of every 100 will suffer disabilities such as hearing loss, brain damage, kidney damage, amputations, nervous system problems, or severe scars from skin grafts.

Meningococcal ACWY vaccine can help prevent meningococcal disease caused by serogroups A, C, W, and Y. A different meningococcal vaccine is available to help protect against serogroup B.

2 Meningococcal ACWY Vaccine

Meningococcal conjugate vaccine (**MenACWY**) is licensed by the Food and Drug Administration (FDA) for protection against serogroups A, C, W, and Y.

Two doses of MenACWY are routinely recommended for adolescents 11 through 18 years old: the first dose at 11 or 12 years old, with a booster dose at age 16. Some adolescents, including those with HIV, should get additional doses. Ask your health care provider for more information.

In addition to routine vaccination for adolescents, MenACWY vaccine is also recommended for certain groups of people:

- People at risk because of a serogroup A, C, W, or Y meningococcal disease outbreak
- People with HIV
- Anyone whose spleen is damaged or has been removed, including people with sickle cell disease
- Anyone with a rare immune system condition called “persistent complement component deficiency”
- Anyone taking a drug called eculizumab (also called Soliris®)
- Microbiologists who routinely work with isolates of *N. meningitidis*
- Anyone traveling to, or living in, a part of the world where meningococcal disease is common, such as parts of Africa
- College freshmen living in dormitories
- U.S. military recruits

Some people need multiple doses for adequate protection. Ask your health care provider about the number and timing of doses, and the need for booster doses.



3

Some people should not get this vaccine

Tell the person who is giving you the vaccine **if you have any severe, life-threatening allergies**. If you have ever had a life-threatening allergic reaction after a previous dose of meningococcal ACWY vaccine, or if you have a severe allergy to any part of this vaccine, you should not get this vaccine. Your provider can tell you about the vaccine's ingredients.

Not much is known about the risks of this vaccine for a pregnant woman or breastfeeding mother. However, pregnancy or breastfeeding are not reasons to avoid MenACWY vaccination. A pregnant or breastfeeding woman should be vaccinated if she is at increased risk of meningococcal disease.

If you have a mild illness, such as a cold, you can probably get the vaccine today. If you are moderately or severely ill, you should probably wait until you recover. Your doctor can advise you.

4

Risks of a vaccine reaction

With any medicine, including vaccines, there is a chance of side effects. These are usually mild and go away on their own within a few days, but serious reactions are also possible.

As many as half of the people who get meningococcal ACWY vaccine have **mild problems** following vaccination, such as redness or soreness where the shot was given. If these problems occur, they usually last for 1 or 2 days.

A small percentage of people who receive the vaccine experience muscle or joint pains.

Problems that could happen after any injected vaccine:

- People sometimes faint after a medical procedure, including vaccination. Sitting or lying down for about 15 minutes can help prevent fainting, and injuries caused by a fall. Tell your doctor if you feel dizzy or lightheaded, or have vision changes.
- Some people get severe pain in the shoulder and have difficulty moving the arm where a shot was given. This happens very rarely.
- Any medication can cause a severe allergic reaction. Such reactions from a vaccine are very rare, estimated at about 1 in a million doses, and would happen within a few minutes to a few hours after the vaccination.

As with any medicine, there is a very remote chance of a vaccine causing a serious injury or death.

The safety of vaccines is always being monitored. For more information, visit: www.cdc.gov/vaccinesafety/

5

What if there is a serious reaction?

What should I look for?

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or unusual behavior.

Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness—usually within a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 and get to the nearest hospital. Otherwise, call your doctor.

Afterward, the reaction should be reported to the "Vaccine Adverse Event Reporting System" (VAERS). Your doctor should file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling 1-800-822-7967.

VAERS does not give medical advice.

6

The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling 1-800-338-2382 or visiting the VICP website at www.hrsa.gov/vaccinecompensation. There is a time limit to file a claim for compensation.

7

How can I learn more?

- Ask your health care provider. He or she can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call 1-800-232-4636 (1-800-CDC-INFO) or
 - Visit CDC's website at www.cdc.gov/vaccines

Vaccine Information Statement (Interim)
**Meningococcal ACWY
Vaccines**



Office use only

08/24/2018 | 42 U.S.C. § 300aa-26

Serogroup B Meningococcal Vaccine (MenB): What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Meningococcal disease is a serious illness caused by a type of bacteria called *Neisseria meningitidis*. It can lead to meningitis (infection of the lining of the brain and spinal cord) and infections of the blood. Meningococcal disease often occurs without warning—even among people who are otherwise healthy.

Meningococcal disease can spread from person to person through close contact (coughing or kissing) or lengthy contact, especially among people living in the same household.

There are at least 12 types of *N. meningitidis*, called “serogroups.” Serogroups A, B, C, W, and Y cause most meningococcal disease.

Anyone can get meningococcal disease but certain people are at increased risk, including:

- Infants younger than one year old
- Adolescents and young adults 16 through 23 years old
- People with certain medical conditions that affect the immune system
- Microbiologists who routinely work with isolates of *N. meningitidis*
- People at risk because of an outbreak in their community

Even when it is treated, meningococcal disease kills 10 to 15 infected people out of 100. And of those who survive, about 10 to 20 out of every 100 will suffer disabilities such as hearing loss, brain damage, kidney damage, amputations, nervous system problems, or severe scars from skin grafts.

Serogroup B meningococcal (MenB) vaccines can help prevent meningococcal disease caused by serogroup B. Other meningococcal vaccines are recommended to help protect against serogroups A, C, W, and Y.

2 Serogroup B Meningococcal Vaccines

Two serogroup B meningococcal vaccines—Bexsero® and Trumenba®—have been licensed by the Food and Drug Administration (FDA).

These vaccines are recommended routinely for people 10 years or older who are at increased risk for serogroup B meningococcal infections, including:

- People at risk because of a serogroup B meningococcal disease outbreak
- Anyone whose spleen is damaged or has been removed
- Anyone with a rare immune system condition called “persistent complement component deficiency”
- Anyone taking a drug called eculizumab (also called Soliris®)
- Microbiologists who routinely work with isolates of *N. meningitidis*

These vaccines may also be given to anyone 16 through 23 years old to provide short term protection against most strains of serogroup B meningococcal disease; 16 through 18 years are the preferred ages for vaccination.

For best protection, more than 1 dose of a serogroup B meningococcal vaccine is needed. The same vaccine must be used for all doses. Ask your health care provider about the number and timing of doses.

3 Some people should not get these vaccines

Tell the person who is giving you the vaccine:

- **If you have any severe, life-threatening allergies.**
If you have ever had a life-threatening allergic reaction after a previous dose of serogroup B meningococcal vaccine, or if you have a severe allergy to any part of this vaccine, you should not get the vaccine. *Tell your health care provider if you have any severe allergies that you know of, including a severe allergy to latex.* He or she can tell you about the vaccine’s ingredients.
- **If you are pregnant or breastfeeding.**
There is not very much information about the potential risks of this vaccine for a pregnant woman or breastfeeding mother. It should be used during pregnancy only if clearly needed.

If you have a mild illness, such as a cold, you can probably get the vaccine today. If you are moderately or severely ill, you should probably wait until you recover. Your doctor can advise you.



4 Risks of a vaccine reaction

With any medicine, including vaccines, there is a chance of reactions. These are usually mild and go away on their own within a few days, but serious reactions are also possible.

More than half of the people who get serogroup B meningococcal vaccine have **mild problems** following vaccination. These reactions can last up to 3 to 7 days, and include:

- Soreness, redness, or swelling where the shot was given
- Tiredness or fatigue
- Headache
- Muscle or joint pain
- Fever or chills
- Nausea or diarrhea

Other problems that could happen after these vaccines:

- People sometimes faint after a medical procedure, including vaccination. Sitting or lying down for about 15 minutes can help prevent fainting and injuries caused by a fall. Tell your provider if you feel dizzy, or have vision changes or ringing in the ears.
- Some people get shoulder pain that can be more severe and longer-lasting than the more routine soreness that can follow injections. This happens very rarely.
- Any medication can cause a severe allergic reaction. Such reactions from a vaccine are very rare, estimated at about 1 in a million doses, and would happen within a few minutes to a few hours after the vaccination.

As with any medicine, there is a very remote chance of a vaccine causing a serious injury or death.

The safety of vaccines is always being monitored. For more information, visit: www.cdc.gov/vaccinesafety/

5 What if there is a serious reaction?

What should I look for?

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or unusual behavior.

Signs of a **severe allergic reaction** can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would usually start a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a **severe allergic reaction** or other emergency that can't wait, call 9-1-1 and get to the nearest hospital. Otherwise, call your clinic.

Afterward the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor should file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS does not give medical advice.

6 The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling **1-800-338-2382** or visiting the VICP website at www.hrsa.gov/vaccinecompensation. There is a time limit to file a claim for compensation.

7 How can I learn more?

- Ask your health care provider. He or she can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines

Vaccine Information Statement Serogroup B Meningococcal Vaccine

08/09/2016

42 U.S.C. § 300aa-26

Office Use Only



Tdap Vaccine

What You Need to Know

(Tetanus,
Diphtheria and
Pertussis)

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Tetanus, diphtheria and pertussis are very serious diseases. Tdap vaccine can protect us from these diseases. And, Tdap vaccine given to pregnant women can protect newborn babies against pertussis.

TETANUS (Lockjaw) is rare in the United States today. It causes painful muscle tightening and stiffness, usually all over the body.

- It can lead to tightening of muscles in the head and neck so you can't open your mouth, swallow, or sometimes even breathe. Tetanus kills about 1 out of 10 people who are infected even after receiving the best medical care.

DIPHThERIA is also rare in the United States today. It can cause a thick coating to form in the back of the throat.

- It can lead to breathing problems, heart failure, paralysis, and death.

PERTUSSIS (Whooping Cough) causes severe coughing spells, which can cause difficulty breathing, vomiting and disturbed sleep.

- It can also lead to weight loss, incontinence, and rib fractures. Up to 2 in 100 adolescents and 5 in 100 adults with pertussis are hospitalized or have complications, which could include pneumonia or death.

These diseases are caused by bacteria. Diphtheria and pertussis are spread from person to person through secretions from coughing or sneezing. Tetanus enters the body through cuts, scratches, or wounds.

Before vaccines, as many as 200,000 cases of diphtheria, 200,000 cases of pertussis, and hundreds of cases of tetanus, were reported in the United States each year. Since vaccination began, reports of cases for tetanus and diphtheria have dropped by about 99% and for pertussis by about 80%.

2 Tdap vaccine

Tdap vaccine can protect adolescents and adults from tetanus, diphtheria, and pertussis. One dose of Tdap is routinely given at age 11 or 12. People who did *not* get Tdap at that age should get it as soon as possible.

Tdap is especially important for healthcare professionals and anyone having close contact with a baby younger than 12 months.

Pregnant women should get a dose of Tdap during **every pregnancy**, to protect the newborn from pertussis. Infants are most at risk for severe, life-threatening complications from pertussis.

Another vaccine, called Td, protects against tetanus and diphtheria, but not pertussis. A Td booster should be given every 10 years. Tdap may be given as one of these boosters if you have never gotten Tdap before. Tdap may also be given after a severe cut or burn to prevent tetanus infection.

Your doctor or the person giving you the vaccine can give you more information.

Tdap may safely be given at the same time as other vaccines.

3 Some people should not get this vaccine

- A person who has ever had a life-threatening allergic reaction after a previous dose of any diphtheria, tetanus or pertussis containing vaccine, OR has a severe allergy to any part of this vaccine, should not get Tdap vaccine. Tell the person giving the vaccine about any severe allergies.
- Anyone who had coma or long repeated seizures within 7 days after a childhood dose of DTP or DTaP, or a previous dose of Tdap, should not get Tdap, unless a cause other than the vaccine was found. They can still get Td.
- Talk to your doctor if you:
 - have seizures or another nervous system problem,
 - had severe pain or swelling after any vaccine containing diphtheria, tetanus or pertussis,
 - ever had a condition called Guillain-Barré Syndrome (GBS),
 - aren't feeling well on the day the shot is scheduled.



4 Risks

With any medicine, including vaccines, there is a chance of side effects. These are usually mild and go away on their own. Serious reactions are also possible but are rare.

Most people who get Tdap vaccine do not have any problems with it.

Mild problems following Tdap (Did not interfere with activities)

- Pain where the shot was given (about 3 in 4 adolescents or 2 in 3 adults)
- Redness or swelling where the shot was given (about 1 person in 5)
- Mild fever of at least 100.4°F (up to about 1 in 25 adolescents or 1 in 100 adults)
- Headache (about 3 or 4 people in 10)
- Tiredness (about 1 person in 3 or 4)
- Nausea, vomiting, diarrhea, stomach ache (up to 1 in 4 adolescents or 1 in 10 adults)
- Chills, sore joints (about 1 person in 10)
- Body aches (about 1 person in 3 or 4)
- Rash, swollen glands (uncommon)

Moderate problems following Tdap (Interfered with activities, but did not require medical attention)

- Pain where the shot was given (up to 1 in 5 or 6)
- Redness or swelling where the shot was given (up to about 1 in 16 adolescents or 1 in 12 adults)
- Fever over 102°F (about 1 in 100 adolescents or 1 in 250 adults)
- Headache (about 1 in 7 adolescents or 1 in 10 adults)
- Nausea, vomiting, diarrhea, stomach ache (up to 1 or 3 people in 100)
- Swelling of the entire arm where the shot was given (up to about 1 in 500).

Severe problems following Tdap (Unable to perform usual activities; required medical attention)

- Swelling, severe pain, bleeding and redness in the arm where the shot was given (rare).

Problems that could happen after any vaccine:

- People sometimes faint after a medical procedure, including vaccination. Sitting or lying down for about 15 minutes can help prevent fainting, and injuries caused by a fall. Tell your doctor if you feel dizzy, or have vision changes or ringing in the ears.
- Some people get severe pain in the shoulder and have difficulty moving the arm where a shot was given. This happens very rarely.
- Any medication can cause a severe allergic reaction. Such reactions from a vaccine are very rare, estimated at fewer than 1 in a million doses, and would happen within a few minutes to a few hours after the vaccination.

As with any medicine, there is a very remote chance of a vaccine causing a serious injury or death.

The safety of vaccines is always being monitored. For more information, visit: www.cdc.gov/vaccinesafety/

5 What if there is a serious problem?

What should I look for?

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or unusual behavior.
- Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would usually start a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 or get the person to the nearest hospital. Otherwise, call your doctor.
- Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor might file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling 1-800-822-7967.

VAERS does not give medical advice.

6 The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling 1-800-338-2382 or visiting the VICP website at www.hrsa.gov/vaccinecompensation. There is a time limit to file a claim for compensation.

7 How can I learn more?

- Ask your doctor. He or she can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call 1-800-232-4636 (1-800-CDC-INFO) or
 - Visit CDC's website at www.cdc.gov/vaccines

Vaccine Information Statement Tdap Vaccine

2/24/2015

42 U.S.C. § 300aa-26



HPV (Human Papillomavirus) Vaccine: What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de Información Sobre Vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

HPV vaccine prevents infection with human papillomavirus (HPV) types that are associated with many cancers, including:

- **cervical cancer** in females,
- **vaginal and vulvar cancers** in females,
- **anal cancer** in females and males,
- **throat cancer** in females and males, and
- **penile cancer** in males.

In addition, HPV vaccine prevents infection with HPV types that cause **genital warts** in both females and males.

In the U.S., about 12,000 women get cervical cancer every year, and about 4,000 women die from it. HPV vaccine can prevent most of these cases of cervical cancer.

Vaccination is not a substitute for cervical cancer screening. This vaccine does not protect against all HPV types that can cause cervical cancer. Women should still get regular Pap tests.

HPV infection usually comes from sexual contact, and most people will become infected at some point in their life. About 14 million Americans, including teens, get infected every year. Most infections will go away on their own and not cause serious problems. But thousands of women and men get cancer and other diseases from HPV.

2 HPV vaccine

HPV vaccine is approved by FDA and is recommended by CDC for both males and females. It is routinely given at 11 or 12 years of age, but it may be given beginning at age 9 years through age 26 years.

Most adolescents 9 through 14 years of age should get HPV vaccine as a two-dose series with the doses separated by 6-12 months. People who start HPV vaccination at 15 years of age and older should get the vaccine as a three-dose series with the second dose given 1-2 months after the first dose and the third dose given 6 months after the first dose. There are several exceptions to these age recommendations. Your health care provider can give you more information.

3 Some people should not get this vaccine

- Anyone who has had a severe (life-threatening) allergic reaction to a dose of HPV vaccine should not get another dose.
- Anyone who has a severe (life threatening) allergy to any component of HPV vaccine should not get the vaccine.

Tell your doctor if you have any severe allergies that you know of, including a severe allergy to yeast.

- HPV vaccine is not recommended for pregnant women. If you learn that you were pregnant when you were vaccinated, there is no reason to expect any problems for you or your baby. Any woman who learns she was pregnant when she got HPV vaccine is encouraged to contact the manufacturer's registry for HPV vaccination during pregnancy at 1-800-986-8999. Women who are breastfeeding may be vaccinated.
- If you have a mild illness, such as a cold, you can probably get the vaccine today. If you are moderately or severely ill, you should probably wait until you recover. Your doctor can advise you.

4 Risks of a vaccine reaction

With any medicine, including vaccines, there is a chance of side effects. These are usually mild and go away on their own, but serious reactions are also possible.

Most people who get HPV vaccine do not have any serious problems with it.

Mild or moderate problems following HPV vaccine:

- Reactions in the arm where the shot was given:
 - Soreness (about 9 people in 10)
 - Redness or swelling (about 1 person in 3)
- Fever:
 - Mild (100°F) (about 1 person in 10)
 - Moderate (102°F) (about 1 person in 65)
- Other problems:
 - Headache (about 1 person in 3)



Problems that could happen after any injected vaccine:

- People sometimes faint after a medical procedure, including vaccination. Sitting or lying down for about 15 minutes can help prevent fainting, and injuries caused by a fall. Tell your doctor if you feel dizzy, or have vision changes or ringing in the ears.
- Some people get severe pain in the shoulder and have difficulty moving the arm where a shot was given. This happens very rarely.
- Any medication can cause a severe allergic reaction. Such reactions from a vaccine are very rare, estimated at about 1 in a million doses, and would happen within a few minutes to a few hours after the vaccination.

As with any medicine, there is a very remote chance of a vaccine causing a serious injury or death.

The safety of vaccines is always being monitored. For more information, visit: www.cdc.gov/vaccinesafety/.

5 What if there is a serious reaction?

What should I look for?

Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or unusual behavior.

Signs of a **severe allergic reaction** can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would usually start a few minutes to a few hours after the vaccination.

What should I do?

If you think it is a **severe allergic reaction** or other emergency that can't wait, call 9-1-1 or get to the nearest hospital. Otherwise, call your doctor.

Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor should file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS does not give medical advice.

6 The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling **1-800-338-2382** or visiting the VICP website at www.hrsa.gov/vaccinecompensation. There is a time limit to file a claim for compensation.

7 How can I learn more?

- Ask your health care provider. He or she can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/hpv

Vaccine Information Statement HPV Vaccine

12/02/2016

42 U.S.C. § 300aa-26

Office Use Only

